

Available Arizona VFC Vaccine & Age Guidelines – Revised June 1, 2011

Vaccines Available from Arizona VFC Program	Age for Vaccine Administration	Special Conditions/Limitations
DT - Diphtheria Tetanus (pediatric)	6 weeks - 6 years	Available upon request to the VFC office as needed.
DTaP - Diphtheria Tetanus acellular Pertussis	6 weeks - 6 years	
DTaP/HepB/IPV - Diphtheria Tetanus acellular Pertussis/Hepatitis B/enhanced Inactivated Polio Vaccine	6 weeks - 6 years- Pediarix	For the first 3 doses of the series only.
DTaP/IPV - Diphtheria Tetanus acellular Pertussis/Inactivated Polio	4 years – 6 years - Kinrix	Used only for the booster dose at age 4-6 years
DTaP/IPV/Hib - Diphtheria Tetanus acellular Pertussis/Inactivated Polio/Haemophilus influenza type B Vaccine	6 weeks – 4 years - Pentacel	For the first 4 doses of DTaP series only
Rotavirus	6 weeks – 32 weeks	Do not give the first dose after 15 weeks of age. Do not administer after 32 weeks of age
PCV13 - Pneumococcal Conjugate Vaccine - 13 valent	6 weeks – 59 months	All children ages of 14 mo. to 59 months, who have received 4 doses of PCV7 should receive 1 dose of PCV13
eIPV - enhanced Inactivated Polio Vaccine	6 weeks - 18 years	
Hib - Haemophilus influenzae type b	6 weeks - 4 years	
Hiberix - Haemophilus influenzae type b	15months-4years only	Can only be used for the “Booster” dose
Hep A - Hepatitis A Vaccine (HAV)	12 months – 18 years	Required for entry to licensed Child Care Centers in Maricopa County only for ages 12 mo. – 5years.
Hep B - Hepatitis B Vaccine (HBV)	Birth - 18 years	
MMR - Measles, Mumps, Rubella	12 months - 18 years	
Varicella - Chicken pox vaccine (Varivax®) *2nd dose now recommended for ages 4 years through 12 years	12 months - 18 years	13 years of age and older require 2 doses separated by 4 to 8 weeks. Less then 13 yrs old should be separated by 3 months
Td - Tetanus diphtheria (adult)	7 years - 18 years	
HPV 4 - Human Papillomavirus - Gardasil	11 years – 18 years-Merck brand only	Now available for Male and Female patients
HPV 2 – Human Papillomavirus –Cervarix	11 years – 18 years-GSK brand only	Available for Females ONLY
MCV4 – Meningococcal Conjugate- Menactra MCV4-Meningococcal Conjugate- Menveo	10 years - 18 years	Administer at age 11-12 yrs-with a booster at age 16 yrs First dose at 13-15 yrs.-administer a booster at 16-18 yrs Administer 2 doses 8 wks apart for ages 2-10 yrs for children with asplenia or complement component deficiency, then 1 dose every 5 yrs thereafter Administer 2 doses 8 wks apart to children with HIV
Tdap - Tetanus Diphtheria acellular Pertussis - Boosterix Tdap - Tetanus Diphtheria acellular Pertussis - Adacel	7 years – 18 years	Use as booster dose - if 5 years or longer have passed since last DTaP/Td – give at age 7 to 9 if incomplete primary series.
PPV23 – Pneumococcal Polysaccharide Vaccine	2 years - 18 years	High risk children only